

Hemangioma of the Median Nerve: A Case Report and Review of Literatures

Piya Kiatisevi, M.D.¹, Supakrit Kijparkorn, M.D.¹, Voranuch Thanakit, M.D.², Koakait Vivitmongkonchai, M.D.³, Bhasanan Sukanthanak, M.D.¹, Piya Assawaboonyadej, M.D.¹

¹ Orthopaedic Oncology Unit, Institute of Orthopaedics, Lerdsin General Hospital Bangkok, Thailand

² Department of Pathology, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

³ Department of Radiology, Faculty of Medicine, Thammasart University, Pathumthani, Thailand

Abstract

Intraneural hemangioma of the median nerve is an extremely rare tumor and only 12 cases have been reported in English literatures. We report a case of 43-year-old man who had an intraneural hemangioma of the median nerve at the right proximal forearm. The patient presented with painful swelling at his right forearm as well as numbness at his index and middle finger for 4 months. Intralesional removal of the mass was achieved by using microscope-assisted epineural and interfascicular dissection. The pain was disappeared after this procedure; however, mild weakness at thumb and index finger flexion was noted. At 12 months follow-up, there was mild weakness noted only at thumb flexion, numbness was much improved and no recurrence was observed.

เสนอโดยนายปิยะ เกียรติเสวี นายแพทย์ชำนาญการพิเศษ โรงพยาบาลเลิดสิน

ในการประชุมวิชาการ 9th Asia Pacific Musculoskeletal Tumor Society Meeting

ซึ่งจัดขึ้นระหว่างวันที่ ๕ - ๑๐ กันยายน ๒๕๕๕ ณ ประเทศมาเลเซีย